

Safe & healthy babies start with safe & healthy pregnancies!



Are you and your baby safe?

HAS YOUR PARTNER EVER?

- Hit, kicked, strangled (choked), or otherwise hurt or threatened you?
- Tried to harm your pregnancy by hitting, kicking, or pushing your stomach?
- Threatened to try to end your pregnancy?
- Blamed you for being pregnant?
- Demanded that you get an abortion?
- Called you names? Insulted you?
- Threatened to kill you or your baby if you leave?
- Threatened to take your baby after birth?
- Tried to control how you care for yourself and your baby?
- Forced you to get pregnant when you didn't want to?
- Refused to use a condom or other contraception when you asked them to?
- Led you to believe they were using contraception when they were not?
- Coerced you, threatened you, and/or used physical violence to keep you from using contraception?
- Tampered with your form of contraception?
- Kept you from seeing a health care provider or seeking medical treatment?
- Tried to keep you from seeing your family and friends?
- Taken your phone?
- Constantly called or texted to see where you were?
- Blamed you for problems with the relationship?
- Accused you of cheating?
- Questioned the paternity of your baby?
- Destroyed your things?
- Hurt or killed your pets?
- Kept you from getting a job or accessing money?
- Tried to sabotage your employment?
- Made you feel like you had to walk on eggshells?
- Done other things that make you feel afraid?

Talk with your healthcare provider. Medical visits are an opportunity to reach out for support.



MARYLAND HEALTH CARE COALITION *Against Domestic Violence*

HealthandDV.org

For more information, scan the QR code on the reverse page.

The Facts About Abuse During Pregnancy



ABUSE CAN BE PHYSICAL, VERBAL, EMOTIONAL, ECONOMIC, PSYCHOLOGICAL, AND/OR SEXUAL.

- Intimate partner violence (IPV) affects millions of people regardless of age, socioeconomic status, race, religion, ethnicity, sexual orientation, or educational background.
- Individuals subjected to IPV may have lifelong health consequences.
- 1 in 3 women globally experiences violence in her lifetime. Many individuals who are pregnant report that abuse began or got worse during pregnancy.
- Abuse is four times more likely to occur if the pregnancy is unwanted.
- Children born to abused mothers are 17% more likely to be born underweight and more than 30% more likely than other children to require intensive care upon birth.
- Health disparities negatively impact maternal and reproductive health outcomes. Abuse from a partner can make these issues worse.

Being free of violence and fear can help you have a safe and healthy pregnancy. Help is available.

National Domestic Violence Hotline (24/7 toll-free support)
1-800-799-SAFE (1-800-799-7233)
TTY 1-800-787-3224



Scan or take a picture of this QR Code to access more information on abuse during pregnancy, possible health risks, strategies to talk to your doctor, and resources for help.



MARYLAND HEALTH CARE COALITION *Against Domestic Violence*

MedChi
The Maryland State Medical Society



HealthandDV.org

1211 Cathedral Street
Baltimore, MD 21201
410-539-0872 or
800-492-1056 x3316

This material was developed through the Local Health Department Preventing Maternal Mortality Funding for Charles County, in partnership with the Center for a Healthy Maryland, the Charles County Department of Health, and survivors of intimate partner violence. The Governor's Office of Crime Prevention and Policy funded the printing of this project under subaward number VOCCG-2025-0056. All points of view in this document are those of the author and do not necessarily represent the official position of any State or Federal agency.